
Choosing the right magnetic beads





Three types of magnetic particles available from Cytiva

| | Sera-Mag™ magnetic beads | SeraSil-Mag™ silica beads | Mag Sepharose™ magnetic beads |
|---------------------------|--|--|---|
| Core material | Carboxyl | Silica | Agarose |
| Surface chemistry options | Streptavidin coated/ Streptavidin blocked Amine-blocked Oligo (dT) coated NeutrAvidin Protein A/G | Silica (OH) | Streptavidin Protein A or G NHS His |
| Size | 1 μm | 400 nm and 700 nm | 37–100 μm |
| Size distribution | Monodisperse | Monodisperse | Polydisperse |
| Surface | Solid but irregular (cauliflower like) | Uniform and spherical | Porous |
| Capacity | Medium | High | High |
| Application focus | Nucleic acid extraction and cleanup Direct conjugation of ligands such as antibodies, enzymes, or oligos mRNA purification Enrichment of target nucleic acid sequences for NGS Size selection of nucleic acids Antibody purification with combined A/G affinities | Traditional nucleic acid extraction using chaotropic salts Existing Sera-Xtracta™ extraction kits use of SeraSil-Mag silica beads to deliver high quality performance | Antibody purification, screening, immunoprecipitation, and pull-down Direct conjugation of ligands such as antibodies, enzymes, and oligos Affinity purification of histidine tagged recombinant proteins |

Comparison of magnetic bead surface chemistries and applications

| Туре | Properties | Applications | Variations | Product | Pack size | Product code |
|-------------------------------|---|--|--|--|-----------|----------------|
| Sera-Mag magnetic beads | | | | | | |
| Sera-Mag carboxylate-modified | Can associate with nucleic acids for direct capture | Conjugation or direct binding applications: | High-speed version available:Sera-Mag SpeedBeads carboxylate-modified | Sera-Mag Carboxylate- Modified [E3] Magnetic Beads | 15 mL | 44152105050250 |
| magnetic beads | Surface suitable for conjugation | Covalent attachmentAffinity purification and pull-down | | | 100 mL | 44152105050350 |
| | through covalent bonding Can capture molecules containing amino groups | | | | 1000 mL | 44152105050450 |
| | | | | Sera-Mag SpeedBead | 15 mL | 65152105050250 |
| | | Nucleic acid isolation and purification | | Carboxylate-Modified [E3] Magnetic Beads | 100 mL | 65152105050350 |
| | | NGS size selection | | | 1000 mL | 65152105050450 |
| | | | | Sera-Mag Carboxylate- | 15 mL | 24152105050250 |
| | | | | Modified [E7] Magnetic Beads | 100 mL | 24152105050350 |
| | | | | | 1000 mL | 24152105050450 |
| | | | | Sera-Mag SpeedBead | 15 mL | 45152105050250 |
| | | | | Carboxylate-Modified [E7] Magnetic Beads | 100 mL | 45152105050350 |
| | | | | | 1000 mL | 45152105050450 |
| Sera-Mag amine-blocked | Surface suitable for conjugation | Conjugation applications, similar to carboxylate-modified beads | High-speed version available:Sera-Mag SpeedBeads amine-blocked | Sera-Mag SpeedBeads Amine-Blocked Magnetic Beads | 1 mL | 19152104011150 |
| magnetic beads | through covalent bonding Non surfactant, non protein | | | | 5 mL | 19152104010150 |
| | Non-surfactant, non-protein- blocked surface | | | | 100 mL | 19152104010350 |
| | Low non-specific binding | | | | | |
| Sera-Mag Oligo(dT)-coated | Hybridizes with mRNA poly-A | mRNA binding applications:mRNA extraction and purificationRT-PCR | | Sera-Mag Oligo (dT) Coated Magnetic Beads | 1 mL | 38152103011150 |
| magnetic beads | tails | | | | 5 mL | 38152103010150 |
| | High colloidal stability | | | | 100 mL | 38152103010350 |
| | | cDNA library construction | | | | |
| | | Subtractive hybridization | | | | |
| | | NGS (RNA sequencing) | | | | |

| Туре | Properties | Applications | Variations | Product | Pack size | Product code |
|-------------------------------|---|--|--|--|-----------|----------------|
| Sera-Mag streptavidin-coated | Binds biotinylated ligands such | Immunoassay and molecular | High-speed version available: Sera-Mag SpeedBeads streptavidin-coated Biotin binding ranges: 2500 to 3500 pmol/mg 3500 to 4500 pmol/mg 4500 to 5500 pmol/mg | Sera-Mag Streptavidin 2500 to 3500 (Low) pmol per mg Magnetic Beads | 1 mL | 30152103011150 |
| magnetic beads | as proteins, nucleic acids, and peptides | biology applications:Sample preparation and | | | 5 mL | 30152103010150 |
| | Covalently bound streptavidin | assay development for genomics and proteomics | | | 100 mL | 30152103010350 |
| | coating Fast reaction kinetics | genomics and proteomics | | Sera-Mag Streptavidin 3500 to 4500 (Med.) pmol per mg Magnetic Beads | 1 mL | 30152104011150 |
| | Low non-specific binding | | | | 5 mL | 30152104010150 |
| | High throughput and precision | | | | 100 mL | 30152104010350 |
| | riigii tiirougiiput una procision | | | Sera-Mag Streptavidin 4500 to 5500 (High) pmol per mg Magnetic Beads | 1 mL | 30152105011150 |
| | | | | | 5 mL | 30152105010150 |
| | | | | | 100 mL | 30152105010350 |
| | | | | Sera-Mag SpeedBeads Streptavidin 3500 to 4500 (Med.) pmol per mg | 1 mL | 66152104011150 |
| | | | | | 5 mL | 66152104010150 |
| | | | | | 100 mL | 66152104010350 |
| Sera-Mag streptavidin-blocked | Binds biotinylated ligands such | High-specificity biotin binding | High-speed version available:Sera-Mag SpeedBeads streptavidin-blocked | Sera-Mag SpeedBeads Streptavidin-Blocked Magnetic Beads | 1 mL | 21152104011150 |
| magnetic beads | as proteins, nucleic acids, and peptides | applications: molecular and immunodiagnostics | | | 5 mL | 21152104010150 |
| | Non-surfactant, non-protein- blocked surface | NGS library preparation | | | 100 mL | 21152104010350 |
| | Lower non-specific binding than streptavidin-coated beads via additional blocking of non-specific binding sites | | | | | |
| Sera-Mag NeutrAvidin-coated | Binds biotinylated ligands such | Alternative to Streptavidin in | High-speed version available: Sera-Mag SpeedBeads NeutrAvidin-coated Biotin binding range: 3500 to 4500 pmol/mg | Sera-Mag SpeedBeads NeutrAvidin-Coated Magnetic Beads | 1 mL | 78152104011150 |
| magnetic beads | as proteins, nucleic acids, and peptides | d immunoassay and molecular biology applications: | | | 5 mL | 78152104010150 |
| | Fast reaction kinetics | Sample preparation and | | | 100 mL | 78152104010350 |
| | Low non-specific binding | assay development for genomics and proteomics | | | | |
| | High throughput and precision | | | | | |

| Туре | Properties | Applications | Variations | Product | Pack size | Product code |
|-------------------------------------|---|---|-------------------------------|----------------------------------|------------|----------------|
| Sera-Mag protein A/G magnetic beads | Binds IgA and IgG proteins | Antibody isolation applications:Affinity purification and pull-down | | Sera-Mag SpeedBeads | 1 mL | 17152104011150 |
| | Coating based on IgA/IgG fusion protein | | | Protein A/G Magnetic Beads | 5 mL | 17152104010150 |
| | Broad binding capabilities | Immunoprecipitation | | | 100 mL | 17152104010350 |
| SeraSil-Mag silica beads | | | | | | |
| SeraSil-Mag silica-coated | Reversibly binds nucleic acids | Applications with low sample | Available in 400 μm or 700 μm | SeraSil-Mag 400 Silica Beads | 5 mL | 29357369 |
| magnetic beads | based on salt concentration Manadisparsa particles with | Augloic acid extraction | particle sizes | | 60 mL | 29357371 |
| | Monodisperse particles with narrow size ranges | Nucleic acid extraction for molecular diagnostics applications such as qPCR | | | 450 mL | 29357372 |
| | | | | | 1000 mL | 29705862 |
| | | | | SeraSil-Mag 700 | 5 mL | 29357373 |
| | | | | Silica Beads | 60 mL | 29357374 |
| | | | | | 450 mL | 29357375 |
| | | | | | 1000 mL | 29705861 |
| Mag Sepharose magnetic beads | | | | | | |
| His-Mag Sepharose Ni magnetic | Highly cross-linked spherical agarose (Sepharose), including magnetite IMAC immobilized with nickel | Small-scale purification and screening of histidine-tagged proteins from different sources | | | 2 × 1 mL | 28967388 |
| beads | | | | | 5 × 1 mL | 28967390 |
| | | | | | 100 mL | 29104065 |
| His Mag Sepharose excel | Strip resistant ligand with strongly bound nickel for immobilized metal ion affinity chromatography (IMAC) | Small-scale capture and purification of histidine-tagged proteins secreted into eukaryotic cell culture supernatants | | His Mag Sepharose excel Beads | 2 × 1 mL | 17371220 |
| magnetic beads | | | | | 5 × 1 mL | 17371221 |
| | | | | | 10 × 1 mL | 17371222 |
| NHS Mag Sepharose magnetic | Coupling of antibodies, aptamers, and proteins through primary amino groups on the molecules to the NHS ligand on NHS Mag Sepharose | Enrichment of target protein for further downstream analyses such as mass spectometry (MS) and electrophoresis techniques | | NHS Mag | 500 μL | 28944009 |
| beads | | | | Sepharose Beads | 4 × 500 μL | 28951380 |

| Туре | Properties | Applications | Variations | Product | Pack size | Product code |
|--|--|--|------------|---------------------------------------|--|----------------------|
| Protein A Mag Sepharose magnetic beads | Maximum binding capacity due to dense coating of Protein A Optimized capacity for enrichment or immunoprecipitation requiring only low amounts of antibody needed | Enrichment of target proteins via immunoprecipitation or pull-down assays Optimised for downstream analyses such as mass spectometry (MS) and electrophoresis techniques | | Protein A Mag Sepharose Beads | 500 μL 4 × 500 μL | 28944006 28951378 |
| Protein A Mag Sepharose Xtra magnetic beads | Maximum binding capacity due to dense coating of Protein A | High capacity small-scale purification and screening of monoclonal and polyclonal antibodies from various species | | Protein A Mag Sepharose Xtra Beads | 2 × 1 mL 5 × 1 mL | 28967056 28967062 |
| Protein G Mag Sepharose magnetic beads | Maximum binding capacity due to dense coating of Protein G Optimized capacity for enrichment or immunoprecipitation requiring only low amounts of antibody needed | Enrichment of target proteins via immunoprecipitation or pull- down assays Optimised for downstream analyses such as mass spectometry (MS) and electrophoresis techniques | | Protein G Mag Sepharose Beads | 500 μL 4 × 500 μL | 28944008 28951379 |
| Protein G Mag Sepharose Xtra magnetic beads | Maximum binding capacity due to dense coating of Protein G | High capacity small-scale purification/screening of monoclonal and polyclonal antibodies from various species | | Protein G Mag Sepharose Xtra Beads | 2 × 1 mL 5 × 1 mL | 28967066 28967070 |
| Streptavidin Mag Sepharose | Utilizes strong interaction between biotin and streptavidin ligand immobilized on magnetic beads | Enrichment of target proteins through immunoprecipitation and purification of biotinylated biomolecules | | Streptavidin Mag Sepharose | 2 × 1 mL 10% slurry 5 × 1 mL 10% slurry | 28985738 28985799 |

cytiva.com

Cytiva and the Drop logo are trademarks of Life Sciences IP Holdings Corporation or an affiliate doing business as Cytiva. Sera-Mag, SeraSil-Mag, Sepharose, and Sera-Xtracta are trademarks of Global Life Sciences Solutions USA LLC or an affiliate doing business as Cytiva.

NeutrAvidin is a trademark of Thermo Fisher Scientific. Any other third-party trademarks are the property of their respective owners.

© 2022–2024 Cytiva

For local office contact information, visit cytiva.com/contact

CY31839-24May24-FL

